

# Spin excitations in stripe-ordered $\text{La}_{2-x}\text{Sr}_x\text{NiO}_4$ ( $x = 0.275$ and $1/3$ )

A.T. Boothroyd<sup>a,\*</sup>, P.G. Freeman<sup>a</sup>, D. Prabhakaran<sup>a</sup>, H. Woo<sup>b,c</sup>, K. Nakajima<sup>d</sup>, J.M. Tranquada<sup>c</sup>, K. Yamada<sup>e</sup>, and C.D. Frost<sup>b</sup>

<sup>a</sup>Department of Physics, University of Oxford, Oxford, OX1 3PU, United Kingdom

<sup>b</sup>ISIS Facility, Rutherford Appleton Laboratory, Chilton, Didcot, OX11 0QX, United Kingdom

<sup>c</sup>Department of Physics, Brookhaven National Laboratory, Upton, New York 11973, USA

<sup>d</sup>Neutron Scattering Laboratory, ISSP, University of Tokyo, Tokai, Ibaraki, Japan

<sup>e</sup>Institute for Chemical Research, Kyoto University, Gokasho, Uji 610-0011, Japan

We report neutron scattering measurements of the spectrum of magnetic excitations in the stripe-ordered phase of  $\text{La}_{2-x}\text{Sr}_x\text{NiO}_4$  ( $x = 0.275$  and  $1/3$ ). The propagating spin excitations follow a similar dispersion relation for the two compositions, but the line widths are broader for  $x = 0.275$  than for  $x = 1/3$ .

It is now well established that charge carriers doped into antiferromagnets tend to segregate into parallel domain walls separating regions with antiferromagnetic (AF) order of the host spins. This so-called stripe ordering is observed in members of the cuprate family of superconductors [1], making it an exciting field for basic research.

The stripe modulation of spin and charge has been characterized extensively in the layered  $\text{La}_{2-x}\text{Sr}_x\text{NiO}_{4+y}$  system [2,3,4], where  $n_h = x + 2y$  is the level of hole doping. Interest is now turning to stripe dynamics. Recently, we reported two separate neutron inelastic scattering studies of the magnetic excitations in the stripe phase of  $\text{La}_{2-x}\text{Sr}_x\text{NiO}_4$  with  $x \simeq 1/3$  [5,6]. In both cases we observed propagating spin-wave modes throughout the Brillouin zone, with a maximum energy of 80–85 meV.

The case of  $x = 1/3$  is special in that the periodicities of the charge and magnetic superlattices are the same and are commensurate with the host lattice. This leads to a very stable stripe order, with a relatively high ordering temperature of 200 K and an in-plane correlation length of several

hundred Å [4]. Here we compare the magnetic excitation spectrum in crystals with  $x = 0.275$  and  $x = 1/3$ . These crystals have similar stripe correlation lengths, but at  $x = 0.275$  the spin and charge periodicities are not the same and are not commensurate with the host lattice.

Neutron inelastic scattering measurements were made on the MAPS chopper spectrometer at the ISIS Facility. The single-crystal samples were mounted in a closed-cycle refrigerator and aligned with the  $c$  axis parallel to the incident beam direction. Scattered neutrons were recorded in large banks of position-sensitive detectors. We analyzed the spectra by making a series of constant-energy slices and projecting the intensity on to the  $(h, k, 0)$  reciprocal lattice plane. The  $c$  component of the scattering vector is unimportant as there are no measurable spin correlations along the  $c$  axis in the range of energies probed.

The diagonal stripe pattern in  $\text{La}_{2-x}\text{Sr}_x\text{NiO}_4$  is twinned, with stripes either parallel to the  $[1, -1, 0]$  or  $[1, 1, 0]$  directions of the tetragonal lattice (cell parameter  $a = 3.8$  Å). These are described by magnetic ordering wave vectors  $(\frac{1}{2}, \frac{1}{2}, l) \pm (\delta, \delta, 0)$  and  $(\frac{1}{2}, \frac{1}{2}, l) \pm (\delta, -\delta, 0)$ , respectively, where  $\delta \simeq n_h/2$ . In our samples we observed an equal population of the twin domains.

\*Corresponding author. Tel.: +44 1865 272376; fax: +44 1865 272400; E-mail address: a.boothroyd1@physics.ox.ac.uk.




Figure 1. Constant-energy scans at  $E = 40$  meV through the magnetic zone centres in stripe-ordered  $\text{La}_{2-x}\text{Sr}_x\text{NiO}_4$  ( $x = 0.275$  and  $1/3$ ). The scans run parallel to the stripe direction. The lines are fits to Gaussian peak profiles.

Figure 1 shows scans at an energy of 40 meV through the magnetic zone centres  $\mathbf{Q}_m = (0.35, 0.35, l)$  for  $x = 0.275$  and  $\mathbf{Q}_m = (1/3, 1/3, l)$  for  $x = 1/3$ , with  $l \simeq 5$  in both cases. The peaks correspond to spin waves propagating parallel to the stripe direction. The positions of the spin-wave peaks are similar for both crystals, but the peaks are much broader for  $x = 0.275$  and only just resolved. At lower energies the widths of the  $x = 0.275$  spin-wave peaks decrease, but are always greater than for  $x = 1/3$ .

By performing Gaussian fits to scans with different energies we determined the two-dimensional spin-wave dispersion. The dispersion perpendicular to the stripe direction is shown in Fig. 2. Reliable values for the peak centres could only be obtained for energies above  $\sim 30$  meV. In this energy range the dispersion for the two crystals is similar, and consistent with that found in Refs. [5,6]. Some differences were observed below 30 meV, and these will be discussed elsewhere.

Perhaps the most interesting result from this study is the fact that the spin excitations of the




Figure 2. Spin wave dispersion perpendicular to the stripe direction in  $\text{La}_{2-x}\text{Sr}_x\text{NiO}_4$ . The points at zero energy are from the magnetic order Bragg peaks.

$x = 0.275$  phase possess a significant intrinsic width which increases with energy, whereas those of  $x = 1/3$  do not. Remember that both compositions exhibit very long range stripe order.

It is possible that the robustness of the  $x = 1/3$  stripes is due to a strong pinning of the commensurate stripes to the lattice. On the other hand, a clear anomaly has been found near 15–20 meV in the spin excitation spectrum of  $x = 1/3$  [5]. This may be due to some kind of resonant spin-charge coupling, and investigation of this energy range in the  $x = 0.275$  sample will be of interest.

## REFERENCES

1. J.M. Tranquada *et al.*, Nature **375** (1995) 561.
2. C.H. Chen, S-W. Cheong, and A.S. Cooper, Phys. Rev. Lett. **71** (1993) 2461.
3. J.M. Tranquada *et al.*, Phys. Rev. Lett. **73** (1994) 1003.
4. H. Yoshizawa *et al.*, Phys. Rev. B **61** (2000) R854.
5. A.T. Boothroyd *et al.*, Phys. Rev. B **67** (2003) 100407(R).
6. P. Bourges *et al.*, Phys. Rev. Lett. **90** (2003) 147202.